



=>

Uploading C:\Program Files\Stnexp\Queries\10799576.str



```
chain nodes :
15 17 23 24 25 26 27 28 29 30 31
ring nodes :
1 2 3 4 5 6 7 8 9 10 11 12 13 14 16 18 19 20 21 22
chain bonds :
7-15 15-16 15-17 23-24 23-27 23-30 24-25 25-26 27-28 28-29 30-31
ring bonds :
1-2 1-6 2-3 3-4 4-5 5-6 5-7 6-10 7-8 8-9 8-11 9-10 9-14 11-12 12-13
13-14 16-18 16-22 18-19 19-20 20-21 21-22
exact/norm bonds :
7-15 15-16 23-24 23-27 23-30 30-31
exact bonds :
15-17 24-25 25-26 27-28 28-29
normalized bonds :
1-2 1-6 2-3 3-4 4-5 5-6 5-7 6-10 7-8 8-9 8-11 9-10 9-14 11-12 12-13
13-14 16-18 16-22 18-19 19-20 20-21 21-22
```

G1:C,O



```
Match level :
1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 6:CLASS 7:CLASS 8:CLASS 9:CLASS
10:CLASS 11:CLASS 12:CLASS 13:CLASS 14:CLASS 15:CLASS 16:CLASS 17:CLASS
18:CLASS 19:CLASS 20:CLASS 21:CLASS 22:CLASS 23:CLASS 24:CLASS 25:CLASS
26:CLASS 27:CLASS 28:CLASS 29:CLASS 30:CLASS 31:CLASS
```

L1 STRUCTURE UPLOADED

=> d l1

L1 HAS NO ANSWERS

L1 STR



Structure attributes must be viewed using STN Express query preparation.

=> s l1

SAMPLE SEARCH INITIATED 17:54:57 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 3 TO ITERATE

100.0% PROCESSED 3 ITERATIONS 2 ANSWERS  
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE \*\*COMPLETE\*\*  
BATCH \*\*COMPLETE\*\*  
PROJECTED ITERATIONS: 3 TO 163  
PROJECTED ANSWERS: 2 TO 124

L2 2 SEA SSS SAM L1

=> s l1 ful

FULL SEARCH INITIATED 17:55:01 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 96 TO ITERATE

100.0% PROCESSED 96 ITERATIONS 66 ANSWERS  
SEARCH TIME: 00.00.01

L3 66 SEA SSS FUL L1

=> file caplus

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

178.36

178.57

FILE 'CAPLUS' ENTERED AT 17:55:04 ON 12 JUN 2008

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is

held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 12 Jun 2008 VOL 148 ISS 24  
FILE LAST UPDATED: 11 Jun 2008 (20080611/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/legal/infopolicy.html>

=> s l3

L4 6 L3

=> d abs fbib hitstr 1-6

L4 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2008 ACS on STN

AB A series of 9-(anilino)acridine and acridine derivs. bearing an alkylating N-mustard residue at C4 of the acridine chromophore were synthesized. The N-mustard pharmacophore was linked to the C4 of the acridine ring with an O-Et (O-C2), O-Pr (O-C3), or O-Bu (O-C4) spacer. It revealed that all newly synthesized compds. were very potent cytotoxic agents against human leukemia and various solid tumors in vitro. These agents did not exhibit cross-resistance against vinblastine-resistant (CCRF-CEM/VBL) or taxol-resistant (CCRF-CEM/taxol) cells. It also showed that these agents were DNA crosslinking agents rather than topoisomerase II inhibitors. Of these agents, two compds. were shown to have potent antitumor activity in nude mice bearing the human breast carcinoma MX-1 xenograft. The therapeutic efficacy of these two agents are comparable to that of taxol.

AN 2006:478646 CAPLUS Full-text  
DN 145:145512

TI Potent Antitumor 9-Anilinoacridines and Acridines Bearing an Alkylating N-Mustard Residue on the Acridine Chromophore: Synthesis and Biological Activity

AU Su, Tsann-Long; Lin, Yi-Wen; Chou, Ting-Chao; Zhang, Xiuguo; Bacherikov, Valeriy A.; Chen, Ching-Huang; Liu, Leroy F.; Tsai, Tsong-Jen  
CS Laboratory of Bioorganic Chemistry, Institute of Biomedical Sciences, Taipei, 115, Taiwan

SO Journal of Medicinal Chemistry (2006), 49(12), 3710-3718  
CODEN: JMCMAR; ISSN: 0022-2623

PB American Chemical Society  
DT Journal

LA English

OS CASREACT 145:145512


IT 898833-68-6P 898833-69-7P 898833-70-0P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
(preparation of

(amino)[[bis(chloroethyl)amino]alkoxy]acridinyl]amino]benzen  
emethanol derivs. and study of their antitumor activity)

RN 898833-68-6 CAPLUS


CN Benzenemethanol, 3-amino-5-[[4-[2-[bis(2-chloroethyl)amino]ethoxy]-9-acridinyl]amino]-, hydrochloride (1:4) (CA INDEX NAME)



●4 HCl

RN 898833-69-7 CAPLUS


CN Benzenemethanol, 3-amino-5-[[4-[4-[bis(2-chloroethyl)amino]butoxy]-9-acridinyl]amino]-, hydrochloride (1:3) (CA INDEX NAME)



●3 HCl

RN 898833-70-0 CAPLUS


CN Carbamic acid, [3-[[4-[2-[bis(2-chloroethyl)amino]ethoxy]-9-acridinyl]amino]-5-(hydroxymethyl)phenyl]-, ethyl ester (9CI) (CA INDEX NAME)



IT 898833-71-1P 898833-72-2P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(preparation of  
 (hydroxy)[[bis(chloroethyl)amino]alkoxy]acridinyl]amino]benz  
 enemethanol derivs. and study of their antitumor activity)  
 RN 898833-71-1 CAPLUS  
 CN Benzenemethanol, 3-[[4-[2-[bis(2-chloroethyl)amino]ethoxy]-9-  
 acridinyl]amino]-5-hydroxy-, hydrochloride (1:2) (CA INDEX NAME)




●2 HCl

RN 898833-72-2 CAPLUS  
 CN Benzenemethanol, 3-[[4-[4-[bis(2-chloroethyl)amino]butoxy]-9-  
 acridinyl]amino]-5-hydroxy-, hydrochloride (2:5) (CA INDEX NAME)



●5/2 HCl

IT 774234-08-1  
 RL: PAC (Pharmacological activity); BIOL (Biological study)  
 (preparation of (phenylamino)acridine derivs. and N-alkyl mustard  
 group-containing acridine derivs. and study of their antitumor activity in  
 comparison with (acridinylamino)[[bis(chloroethyl)amino]ethoxy]benzenem  
 ethanol)  
 RN 774234-08-1 CAPLUS  
 CN Benzenemethanol, 3-(9-acridinylamino)-5-[2-[bis(2-  
 chloroethyl)amino]ethoxy]- (CA INDEX NAME)




IT 898833-73-3P 898833-74-4P 898833-75-5P  
 898833-76-6P 898833-77-7P 898833-78-8P  
 898833-79-9P 898833-80-2P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL  
 (Biological study); PREP (Preparation)  
 (preparation of [[bis(chloroethyl)amino]alkoxy]acridinyl]benzenediamine  
 derivs. and study of their antitumor activity)

RN 898833-73-3 CAPLUS


CN 1,3-Benzenediamine, N1-[4-[2-[bis(2-chloroethyl)amino]ethoxy]-9-acridinyl]-  
 5-methyl-, hydrochloride (1:7) (CA INDEX NAME)



●7 HCl

RN 898833-74-4 CAPLUS


CN 1,3-Benzenediamine, N1-[4-[4-[bis(2-chloroethyl)amino]butoxy]-9-acridinyl]-  
 5-methyl-, hydrochloride (1:6) (CA INDEX NAME)



●6 HCl

RN 898833-75-5 CAPLUS


CN 1,3-Benzenediamine, N1-[4-[2-[bis(2-chloroethyl)amino]ethoxy]-9-acridinyl]-5-methoxy-, hydrochloride (1:4) (CA INDEX NAME)



●4 HCl

RN 898833-76-6 CAPLUS


CN 1,3-Benzenediamine, N1-[4-[4-[bis(2-chloroethyl)amino]butoxy]-9-acridinyl]-5-methoxy-, hydrochloride (1:4) (CA INDEX NAME)



●4 HCl

RN 898833-77-7 CAPLUS


CN 1,3-Benzenediamine, N3-[4-[2-[bis(2-chloroethyl)amino]ethoxy]-9-acridinyl]-4-methyl-, hydrochloride (1:4) (CA INDEX NAME)



●4 HCl

RN 898833-78-8 CAPLUS


CN 1,3-Benzenediamine, N3-[4-[4-[bis(2-chloroethyl)amino]butoxy]-9-acridinyl]-4-methyl-, hydrochloride (1:4) (CA INDEX NAME)



●4 HCl

RN 898833-79-9 CAPLUS

CN 1,3-Benzenediamine, N1-[4-[2-bis(2-chloroethyl)amino]ethoxy]-9-acridinyl]-4-methoxy-, hydrochloride (1:4) (CA INDEX NAME)




●4 HCl

RN 898833-80-2 CAPLUS

CN 1,3-Benzenediamine, N1-[4-[4-bis(2-chloroethyl)amino]butoxy]-9-acridinyl]-




4-methoxy-, hydrochloride (10:67) (CA INDEX NAME)



● 67/10 HCl

RE.CNT 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2008 ACS on SIN  
GI



I

AB A series of N-mustard derivs. of 9-anilinoacridine was synthesized for antitumor and structure-activity relationship studies. The alkylating N-mustard residue was linked to the C-3' or C-4' position of the anilino ring with an O-ethylene (O-C2), O-butylene (O-C4), and methylene (C1) spacer. All of the new N-mustard derivs. exhibited significant cytotoxicity in inhibiting human lymphoblastic leukemic cells (CCRF-CEM) in culture. Of these agents, (3-(acridin-9-ylamino)-5-(2-[bis (2-chloroethyl)amino]ethoxy)phenyl)methanol (I) was subjected to antitumor studies, resulting in an approx. 100-fold more potent effect than its parent analog 3-(9-acridinylamino)-5-hydroxymethylaniline (AHMA) in inhibiting the growth of human lymphoblastic leukemic cells (CCRF-CEM) in vitro. This agent did not exhibit cross-resistance against vinblastine-resistant (CCRF-CEM/VBL) or Taxol-resistant (CCRF-CEM/Taxol) cells. Remarkably, the therapeutic effect of I at a dose as

low as one tenth of the Taxol therapeutic dose [i.e., 1-2 mg/kg (Q3D + 7) or 3 mg/kg (Q4D + 5); i.v. injection] on nude mice bearing human breast carcinoma MX-1 xenografts resulted in complete tumor remission in two out of three mice. Furthermore, I yielded xenograft tumor suppression of 81-96% using human T-cell acute lymphoblastic leukemia CCRF-CEM, colon carcinoma HCT-116, and ovarian adenocarcinoma SK-OV-3 tumor models.

AN 2005:464998 CAPLUS [Full-text](#)

DN 143:125829

TI Potent antitumor 9-anilinoacridines bearing an alkylating N-mustard residue on the anilino ring: synthesis and biological activity

AU Bacherikov, Valeriy A.; Chou, Ting-Chao; Dong, Hua-Jin; Zhang, Xinguo; Chen, Ching-Huang; Lin, Yi-Wen; Tsai, Tsong-Jen; Lee, Rong-Zau; Liu, Leroy F.; Su, Tsann-Long

CS Laboratory of Bioorganic Chemistry, Institute of Biomedical Sciences, Academia Sinica, Taipei, 115, Taiwan

SO Bioorganic & Medicinal Chemistry (2005), 13(12), 3993-4006

CODEN: BMECEP; ISSN: 0968-0896

PB Elsevier Ltd.

DT Journal

LA English

OS CASREACT 143:125829

IT 774234-08-1P 858136-02-4P 858136-03-5P

858136-04-6P 858136-05-7P 858136-06-8P

858136-07-9P 858136-08-0P 858136-09-1P

858136-10-4P 858136-11-5P 858136-12-6P

858136-13-7P 858136-14-8P 858136-15-9P


858136-16-0P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(antitumor 9-anilinoacridines bearing an alkylating N-mustard residue on the anilino ring)


RN 774234-08-1 CAPLUS

CN Benzenemethanol, 3-(9-acridinylamino)-5-[2-[bis(2-chloroethyl)amino]ethoxy]- (CA INDEX NAME)



RN 858136-02-4 CAPLUS


CN 9-Acridinamine, N-[3-[2-[bis(2-chloroethyl)amino]ethoxy]phenyl]-, hydrochloride (1:1) (CA INDEX NAME)



● HCl

RN 858136-03-5 CAPLUS


CN 9-Acridinamine, N-[3-[4-[bis(2-chloroethyl)amino]butoxy]phenyl]-, hydrochloride (1:2) (CA INDEX NAME)



●2 HCl

RN 858136-04-6 CAPLUS

CN 9-Acridinamine, N-[4-[2-[bis(2-chloroethyl)amino]ethoxy]phenyl]-, hydrochloride (1:2) (CA INDEX NAME)




●2 HCl

RN 858136-05-7 CAPLUS

CN 9-Acridinamine, N-[4-[4-[bis(2-chloroethyl)amino]butoxy]phenyl]-,


hydrochloride (1:3) (CA INDEX NAME)



●3 HCl

RN 858136-06-8 CAPLUS


CN 4-Acridinecarboxamide, 9-[[3-[2-[bis(2-chloroethyl)amino]ethoxy]phenyl]amino]-N-[2-(dimethylamino)ethyl]-5-methyl-, hydrochloride (1:4) (CA INDEX NAME)



●4 HCl

RN 858136-07-9 CAPLUS


CN 4-Acridinecarboxamide, 9-[[3-[4-[bis(2-chloroethyl)amino]butoxy]phenyl]amino]-N-[2-(dimethylamino)ethyl]-5-methyl-, hydrochloride (1:3) (CA INDEX NAME)



●3 HCl

RN 858136-08-0 CAPLUS  
 CN 4-Acridinecarboxamide, 9-[[4-[2-[bis(2-chloroethyl)amino]ethoxy]phenyl]amino]-N-[2-(dimethylamino)ethyl]-5-methyl-, hydrochloride (2:5) (CA INDEX NAME)


PAGE 1-A



PAGE 2-A


●5/2 HCl

RN 858136-09-1 CAPLUS  
 CN 4-Acridinecarboxamide, 9-[[4-[4-bis(2-chloroethyl)amino]butoxy]phenyl]amino]-N-[2-(dimethylamino)ethyl]-5-methyl-, hydrochloride (2:3) (CA INDEX NAME)




● 3/2 HCl

RN 858136-10-4 CAPLUS  
 CN 9-Acridinamine, N-[4-[2-[bis(2-chloroethyl)amino]ethoxy]-3-methylphenyl]-,  
 hydrochloride (2:3) (CA INDEX NAME)



● 3/2 HCl


RN 858136-11-5 CAPLUS  
 CN 9-Acridinamine, N-[4-[2-[bis(2-chloroethyl)amino]ethoxy]-2-methylphenyl]-,  
 hydrochloride (2:3) (CA INDEX NAME)



● 3/2 HCl

RN 858136-12-6 CAPLUS


CN 9-Acridinamine, N-[3-[2-[bis(2-chloroethyl)amino]ethoxy]-4-methoxyphenyl]-, hydrochloride (4:5) (CA INDEX NAME)



● 5/4 HCl

RN 858136-13-7 CAPLUS

CN 9-Acridinamine, N-[3-[[bis(2-chloroethyl)amino]methyl]phenyl]-, hydrochloride (1:2) (CA INDEX NAME)




● 2 HCl

RN 858136-14-8 CAPLUS

CN 9-Acridinamine, N-[4-[[bis(2-chloroethyl)amino]methyl]phenyl]-,


hydrochloride (1:2) (CA INDEX NAME)



● 2 HCl

RN 858136-15-9 CAPLUS

CN 4-Acridinecarboxamide, 9-[[3-[[bis(2-chloroethyl)amino]methyl]phenyl]amino]-N-[2-(dimethylamino)ethyl]-5-methyl-, hydrochloride (1:3) (CA INDEX NAME)




● 3 HCl

RN 858136-16-0 CAPLUS


CN 4-Acridinecarboxamide, 9-[[4-[[bis(2-chloroethyl)amino]methyl]phenyl]amino]-N-[2-(dimethylamino)ethyl]-5-methyl-, hydrochloride (1:3) (CA INDEX NAME)





RE.CNT 43 THERE ARE 43 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 3 OF 6 CAPLUS COPYRIGHT 2008 ACS on STN  
GI



I

AB A series of 9-anilinoacridine N-mustard derivs., in which the alkylating N-mustard residue was linked to the C-3' or C-4' position of the anilino ring with an O-ethylene spacer, was synthesized and evaluated for cytotoxicity against human lymphoblastic leukemic cells (CCRF-CEM) in culture. The results showed that all of the new compds. exhibited potent cytotoxicity with IC50 values ranging from 0.002 to 0.7  $\mu$ M, which were as potent or significantly more potent than 3-(9-acridinylamino)-5-hydroxymethylaniline (AHMA). Compound I did not exhibit cross-resistance against both vinblastine-resistant (CCRF-CEM/VBL) and taxol-resistant (CCRF-CEM/taxol) cells. Addnl., compound I demonstrated potent antitumor effect in nude mice bearing human breast carcinoma MX-1 xenografts, resulting in complete tumor remission in two out of three mice at the maximal dose of 1-2mg/kg (Q3D+7) or 3mg/kg (Q4D+5) via i.v. injection.


AN 2004:689256 CAPLUS Full-text

DN 141:342864


TI Potent antitumor N-mustard derivatives of 9-anilinoacridine, synthesis and antitumor evaluation

AU Bacherikov, Valeriy A.; Chou, Ting-Chao; Dong, Hua-Jin; Chen, Ching-Huang;


Lin, Yi-Wen; Tsai, Tsong-Jen; Su, Tsann-Long  
 CS Laboratory of Bioorganic Chemistry, Institute of Biomedical Sciences,  
 Academia Sinica, Taipei, 115, Taiwan  
 SO Bioorganic & Medicinal Chemistry Letters (2004), 14(18), 4719-4722  
 CODEN: BMCLE8; ISSN: 0960-894X  
 PB Elsevier B.V.  
 DT Journal  
 LA English  
 OS CASREACT 141:342864  
 IT 774234-06-1P 774234-11-6P 774234-12-7P  
 774234-13-9P 774234-14-9P  
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU  
 (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES  
 (Uses)  
 (antitumor N-mustard derivs. of acridine)  
 RN 774234-08-1 CAPLUS  
 CN Benzenemethanol, 3-(9-acridinylamino)-5-[2-[bis(2-  
 chloroethyl)amino]ethoxy]- (CA INDEX NAME)




RN 774234-11-6 CAPLUS  
 CN 9-Acridinamine, N-[3-[2-[bis(2-chloroethyl)amino]ethoxy]phenyl]- (CA  
 INDEX NAME)




RN 774234-12-7 CAPLUS  
 CN 4-Acridinecarboxamide, 9-[[3-[2-[bis(2-chloroethyl)amino]ethoxy]phenyl]ami  
 no]-N-[2-(dimethylamino)ethyl]-5-methyl- (CA INDEX NAME)



RN 774234-13-8 CAPLUS  
 CN 9-Acridinamine, N-[4-[2-[bis(2-chloroethyl)amino]ethoxy]phenyl]- (CA INDEX NAME)




RN 774234-14-9 CAPLUS  
 CN 4-Acridinecarboxamide, 9-[4-[2-[bis(2-chloroethyl)amino]ethoxy]phenyl]amino]-N-[2-(dimethylamino)ethyl]-5-methyl- (CA INDEX NAME)



RE.CNT 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2008 ACS on STN



AB A possible mode of action involving electron transfer is advanced for the 9-anilinoacridines [I; R1 = H, OMe; R2 and R4 = H, Cl; R3 = H, Me; R5 = H, OH, OEt, CO2H; R6 = OH, CH2Net2, CH2N(CH2)4, CH2N(CH2CH2Cl)2, CH2N[(CH2)7Me]2]. The mechanism entails formation of toxic oxy radicals which destroy the neoplasm. Cyclic voltammetry was performed on iminium type ions derived by protonation of the acridines. Redns. were generally reversible with potentials of about -0.60 V. Involvement of quinoidal metabolites is also a possibility. The relationship of electrochem. behavior to structure and physiol. activity is addressed.

AN 1987:628450 CAPLUS [Full-text](#)

DN 107:228450

OREF 107:36495a,36498a

TI Electron transfer-oxy radical mechanism for anticancer agents: 9-anilinoacridines

AU Kovacic, P.; Ames, J. R.; Ryan, M. D.

CS Dep. Chem., Univ. Wisconsin, Milwaukee, WI, 53201, USA

SO Anti-Cancer Drug Design (1987), 2(1), 37-46

CODEN: ACDDEA; ISSN: 0266-9536

DT Journal

LA English


IT 111393-51-2

RL: BIOL (Biological study)

(electron transfer-oxy radical mechanisms for antitumor, structure in relation to)


RN 111393-51-2 CAPLUS

CN Phenol, 2-[[bis(2-chloroethyl)amino]methyl]-4-[(6-chloro-2-methoxy-9-acridinyl)amino]-, dihydrochloride (9CI) (CA INDEX NAME)




●2 HCl


L4 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2008 ACS on STN  
 AB cf. CA 54, 6954d. This indexed compilation contains information concerning  
 1816 compds. tested.  
 AN 1960:131417 CAPLUS [Full-text](#)  
 DN 54:131417  
 OREF 54:25245g  
 TI Cancer chemotherapy screening data. VII  
 AU Leiter, Joseph  
 SO Cancer Research (1960), 20(No. 7,Pt. 2), 471-684  
 CODEN: CNREA8; ISSN: 0008-5472  
 DT Journal  
 LA Unavailable  
 IT 102892-27-3, o-Cresol,  $\alpha$ -[bis(2-chloroethyl)amino]-4-[(6-chloro-2-methoxy-9-acridinyl)amino]- 115918-36-0, o-Cresol,  $\alpha$ -[bis(2-chloroethyl)amino]-4-[(6-chloro-2-phenyl-9-acridinyl)amino]- 121814-84-4, o-Cresol, 4-benz[c]acridin-7-ylamino- $\alpha$ -[bis(2-chloroethyl)amino]- (as cancer inhibitor)  
 RN 102892-27-3 CAPLUS  
 CN Phenol, 2-[[bis(2-chloroethyl)amino]methyl]-4-[(6-chloro-2-methoxy-9-acridinyl)amino]- (CA INDEX NAME)



RN 115918-36-0 CAPLUS  
 CN o-Cresol,  $\alpha$ -[bis(2-chloroethyl)amino]-4-[(6-chloro-2-phenyl-9-acridinyl)amino]- (6CI) (CA INDEX NAME)



RN 121814-84-4 CAPLUS  
 CN o-Cresol, 4-benz[c]acridin-7-ylamino- $\alpha$ -[bis(2-chloroethyl)amino]- (6CI) (CA INDEX NAME)



L4 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2008 ACS on STN

AB Diaminocresols and salts thereof, useful as agricultural fungicides against *Alternaria solani*, are prepared by converting the alcoholic hydroxyl groups in a substituted cresol to the chloride with  $\text{SOCl}_2$ . 2,2'-(5-(7-Chloro-4-quinolylamino)-2-hydroxybenzylimino)diethanol (I) is prepared as follows: 30 g. paraformaldehyde, 105 g. diethanolamine, and 100 ml. EtOH is boiled to a clear solution to which is added a hot solution of 151 g. 4'-hydroxyacetanilide in 100 ml. EtOH. The mixture is refluxed 3 hrs., 500 ml. 1:1-HCl added, refluxed 2 hrs., the residue dried thoroughly and treated with 198 g. 4,7-dichloroquinoline and 400 g. PhOH and heated 3 hrs. at  $130-140^\circ$ . The melt is acidified with alc. HCl, precipitated with five vols. of acetone to a gum which is triturated with Et<sub>2</sub>O to a granular solid. Acetone containing a slight excess of  $\text{NH}_4\text{OH}$  extracts the base. The acetone solution is decolorized with C, concentrated to 2 l. and diluted with 1.5 vols. hot H<sub>2</sub>O to give yellow I, m.  $193-4^\circ$  (50% EtOH). 4,7-Dichloroquinoline (81.5 g.), 68.8 g. p-methylaminophenol sulfate, and 150 g. PhOH is heated 2 hrs. at  $140^\circ$ , then heated in succession with a few drops of alc. HCl, one volume acetone and 5 vols. Et<sub>2</sub>O. The precipitated red oil is triturated with anhydrous Et<sub>2</sub>O and with excess saturated  $\text{NaHCO}_3$  solution. The precipitate collected by filtration, washed with ether, saturated  $\text{NaHCO}_3$  and water, is dried and digested with 400 ml. 95% EtOH to give 4-[(7-chloro-4-quinolyl)methylamino]phenol (II), m.  $256-8^\circ$  (EtOH). A mixture of 15.7 g. diethanolamine, 4.5 g. paraformaldehyde, and 100 ml. ProH is heated to a solution which is added to a suspension of 27.5 g. II in 250 ml. ProH and refluxed 4 hrs. The mixture is repeatedly filtered hot after boiling with fresh amts. of diethanolamine-paraformaldehyde solution. The combined filtrate is concentrated in vacuo to a sirup which is made alkaline with NaOH and extracted with  $\text{CHCl}_3$ . The  $\text{CHCl}_3$  extract, washed with 5% NaOH solution, water, is evaporated in vacuo to a gum. The alc. solution of the residue is treated with C and excess alc. HCl and poured into 2 l. Et<sub>2</sub>O to give hygroscopic, yellow 2,2'-(5-[(7-chloro-4-quinolyl)amino]-2-hydroxybenzylimino)diethanol-2HCl.3/4H<sub>2</sub>O (III) of indefinite m.p. Hydrated 2,2'-(5-(benz[c]acridin-7-ylamino)-2-hydroxybenzylimino)diethanol-2HCl (IV), m.  $105-10^\circ$  (decomposition), is prepared likewise from 29.9 g. 2,2'-(5-amino-2-hydroxybenzylimino)diethanol-2HCl, 29.9 g. 7-chlorobenz[c]acridine and 75 g. phenol. I (5.8 g.) is added in portions to 20 ml.  $\text{SOCl}_2$  with stirring. After 16 hrs. at room temperature excess  $\text{SOCl}_2$  is removed by decantation, the brown oily residue is taken up in absolute EtOH, treated with C, filtered and evaporated to about 50 ml. The cooled alc. solution is poured with vigorous stirring into 500 ml. anhydrous Et<sub>2</sub>O to give  $\alpha$ -[bis(2-chloroethyl)amino]-4-(7-chloro-4-quinolylamino)-o-cresol-2HCl.1.5H<sub>2</sub>O, m.  $120^\circ$  (decomposition). In other examples the volume  $\text{SOCl}_2$ , weight of starting heterocyclic alc., and the product and m.p. obtained are: 30 ml., 4 g. 2,2'-(5-(benzo[h]quinolin-4-

ylamino)-2- hydroxybenzylimino]diethanol, 4-(benzo[h]quinolin-4-ylamino)- $\alpha$ -[bis(2-chloroethyl)amino]-o-cresol-2HCl, -; 65 ml., 8.0 g. 2,2'-[5-(benzo[f]quinolin-1-ylamino)-2-hydroxybenzylimino]diethanol, 4-(benzo[f]quinolin-1-ylamino)- $\alpha$ -[bis(2-chloroethyl)amino]-o-cresol-2HCl, -; 20 ml., 4.7 g. III,  $\alpha$ -[bis(2-chloroethyl)amino]-4-[(7-chloro-4-quinolyl)methylamino]-o-cresol-2HCl, -; 200 ml., 36.8 g. 2,2'-[5-(6-chloro-2-methoxyacridin-9-ylamino)-2- hydroxybenzylimino]diethanol,  $\alpha$ -[bis(2-chloroethyl)amino]-4-(6- chloro-2-methoxy-9-acridinylamino)-o-cresol-2HCl, 195-6° (decomposition); 100 ml., 5.3 g. 2,2'-[5-(benz[b]acridin-12-ylamino)-2- hydroxybenzylimino]diethanol-2HCl, 4-(benz[b]acridin-12-ylamino)- $\alpha$ -[bis(2-chloroethyl)amino]-o-cresol-2HCl, -; 150 ml., 10.6 g. IV, 4-(benz[c]-acridin-7-ylamino)- $\alpha$ -[bis(2-chloroethyl)amino]-o-cresol-2HCl, -; 400 ml., 21.2 g. 2,2'-[5-(benz[a]acridin-12-ylamino)-2- hydroxybenzylimino]diethanol-2HCl, 4-(benz[a]acridin-12-ylamino)- $\alpha$ -[bis(2-chloroethyl)amino]-o-cresol-2HCl, -.

AN 1959:89542 CAPLUS Full-text

DN 53:89542

OREF 53:16165f-i,16166a-e

TI Benzoquinolylamino-2-[bis( $\beta$ -chloroethyl)aminomethyl]phenols

IN Elslager, E. F.; Tendick, F. H.


PA Parke, Davis & Co.

DT Patent

LA Unavailable

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2883382		19590421	US 1957-680928	19570829
IT	111393-51-2P, o-Cresol, $\alpha$ -[bis(2-chloroethyl)amino]-4-[(6-chloro-2-methoxy-9-acridinyl)amino]-, dihydrochloride 121814-85-5P, o-Cresol, 4-benz[b]acridin-12-ylamino- $\alpha$ -[bis(2-chloroethyl)amino]-, dihydrochloride 122021-13-0P, o-Cresol, 4-benz[c]acridin-7-ylamino- $\alpha$ -[bis(2-chloroethyl)amino]-, dihydrochloride 859924-40-6P, o-Cresol, 4-benz[a]acridin-12-ylamino- $\alpha$ -[bis(2-chloroethyl)amino]- RL: PREP (Preparation) (preparation of)				
RN	111393-51-2 CAPLUS				
CN	Phenol, 2-[[bis(2-chloroethyl)amino]methyl]-4-[(6-chloro-2-methoxy-9-acridinyl)amino]-, dihydrochloride (9CI) (CA INDEX NAME)				




● 2 HCl

RN 121814-85-5 CAPLUS

CN o-Cresol, 4-benz[b]acridin-12-ylamino- $\alpha$ -[bis(2-chloroethyl)amino]-,


dihydrochloride (6CI) (CA INDEX NAME)



● 2 HCl

RN 122021-13-0 CAPLUS

CN o-Cresol, 4-benz[c]acridin-7-ylamino- $\alpha$ -[bis(2-chloroethyl)amino]-, dihydrochloride (6CI) (CA INDEX NAME)



● 2 HCl

RN 859924-40-6 CAPLUS

CN Phenol, 4-(benz[a]acridin-12-ylamino)-2-[[bis(2-chloroethyl)amino]methyl]- (CA INDEX NAME)

